

VIAGRA can help a man with ED get and keep an erection when he becomes sexually stimulated. But it's not right for everyone. Here are some other things you should know:

DO NOT TAKE VIAGRA IF YOU:

- take any other medicines called nitrates, often prescribed for chest pain, as this may cause a sudden, unsafe drop in blood pressure.
- are allergic to sildenafil, as contained in VIAGRA and REVATIO, or any of the ingredients in VIAGRA.

Discuss your health with your doctor to ensure that you are healthy enough for sex. If you experience chest pain, dizziness, or nausea during sex, seek immediate medical help.

Talk to your doctor.

Discuss your general health and ED symptoms with your doctor to see if VIAGRA is right for you.

- Patients should always ask their doctors for medical advice about adverse events.
- You may report an adverse event related to Pfizer products by calling 1-800-438-1985 (U.S. only). If you prefer, you may contact the U.S. Food and Drug Administration (FDA) directly. The FDA has established a reporting service known as MedWatch where healthcare professionals and consumers can report serious problems they suspect may be associated with the drugs and medical devices they prescribe, dispense, or use. Visit [MedWatch](#) or call 1-800-FDA-1088.
- **Terms and conditions apply.** This is not health insurance. This will only be accepted at participating pharmacies. No membership fees.
- Pfizer Inc | PO Box 29387 | Mission, KS 66201-9618 | 1-855-284-2472

Scroll for Important Safety Information and Indication.

IMPORTANT SAFETY INFORMATION

Do not take VIAGRA if you:

- take any other medicines called nitrates, often prescribed for chest pain, as this may cause a sudden, unsafe drop in blood pressure.
- are allergic to sildenafil, as contained in VIAGRA and REVATIO, or any of the ingredients in VIAGRA.

Discuss your health with your doctor to ensure that you are healthy enough for sex. If you experience chest pain, dizziness, or nausea during sex, seek immediate medical help.

VIAGRA can cause serious side

effects. Rarely reported side effects include:

- **an erection that will not go away (priapism).** If you have an erection that lasts more than 4 hours, get medical help right away. If it is not treated right away, priapism can permanently damage your penis.
- **sudden vision loss in one or both eyes.** Sudden vision loss in one or both eyes can be a sign of a serious eye problem called non-arteritic anterior ischemic optic neuropathy (NAION). Stop taking VIAGRA and call your healthcare provider right away if you have any sudden vision loss.
- **sudden hearing decrease or hearing loss.** Some people may also have ringing in their ears (tinnitus) or dizziness. If you have these symptoms, stop taking VIAGRA and contact a doctor right away.

Before you take VIAGRA, tell your healthcare provider if you:

- have or have had heart problems such as a heart attack, irregular heartbeat, angina, chest pain, narrowing of the aortic valve, or heart failure
- have had heart surgery within the last 6 months
- have had a stroke
- have low blood pressure, or high blood pressure that is not controlled
- have a deformed penis shape
- have had an erection that lasted for more than 4 hours
- have problems with your blood cells such as sickle cell anemia, multiple myeloma, or leukemia
- have retinitis pigmentosa, a rare genetic (runs in families) eye disease
- have ever had severe vision loss, including an eye problem called NAION

- have bleeding problems
- have or have had stomach ulcers
- have liver problems
- have kidney problems or are having kidney dialysis
- have any other medical conditions

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

VIAGRA may affect the way other medicines work, and other medicines may affect the way VIAGRA works, causing side effects. Especially tell your healthcare provider if you take any of the following:

- medicines called nitrates
- medicines called alpha-blockers such as Hytrin (terazosin HCl), Flomax (tamsulosin HCl), Cardura (doxazosin mesylate), Minipress (prazosin HCl), Uroxatral (alfuzosin HCl), Jalyn (dutasteride and tamsulosin HCl), or Rapaflo (silodosin). Alpha-blockers are sometimes prescribed for prostate problems or high blood pressure. In some patients, the use of VIAGRA with alpha-blockers can lead to a drop in blood pressure or to fainting
- medicines called HIV protease inhibitors, such as ritonavir (Norvir), indinavir sulfate (Crixivan), saquinavir (Fortovase or Invirase), or atazanavir sulfate (Reyataz)
- some types of oral antifungal medicines, such as ketoconazole (Nizoral) and itraconazole (Sporanox)
- some types of antibiotics, such as clarithromycin (Biaxin), telithromycin (Ketek), or erythromycin
- other medicines that treat high blood pressure
- other medicines or treatments for ED

- VIAGRA contains sildenafil, which is the same medicine found in another drug called REVATIO. REVATIO is used to treat a rare disease called pulmonary arterial hypertension (PAH). VIAGRA should not be used with REVATIO or with other PAH treatments containing sildenafil or any other PDE5 inhibitors (such as Adcirca [tadalafil])

VIAGRA does not protect against sexually transmitted diseases, including HIV.

The most common side effects of VIAGRA: headache; flushing; upset stomach; abnormal vision, such as changes in color vision (such as having a blue color tinge) and blurred vision; stuffy or runny nose; back pain; muscle pain; nausea; dizziness; rash.

INDICATION

VIAGRA (sildenafil citrate) is prescription medicine used to treat erectile dysfunction (ED).

VIAGRA is not for women or children.

Please see full prescribing information for VIAGRA (25 mg, 50 mg, 100 mg) tablets.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088. By clicking on this link you will be leaving VIAGRA.com. This website is neither owned nor controlled by Pfizer. Pfizer is not responsible for the content or services of this site.

- [VIAGRA and ED](#)
- [BUY REAL VIAGRA](#)

- [TAKING VIAGRA](#)
- [COMMON QUESTIONS](#)

- [Site map](#)
- [For partners](#)
- [Contact us](#)
- [Terms of use](#)
- [Privacy Policy](#)



© 2001-2014 Pfizer Inc. All rights reserved.

The blue diamond tablet shape is a registered trademark of Pfizer Inc.

[REQUEST A BROCHURE](#)

VIAGRA (sildenafil citrate), REVATIO (sildenafil), Cardura (doxazosin mesylate), and Minipress (prazosin HCl) are registered trademarks of Pfizer Inc. Other brands listed are trademarks of their respective owners and are not affiliated with nor endorse Pfizer Inc.

The health information contained herein is provided for educational purposes only and is not intended to replace discussions with a healthcare provider. All decisions regarding patient care must be made with a healthcare provider, considering the unique characteristics of the patient.

The product information provided in this site is intended only for residents of the United States. The products discussed herein may have different product labeling in different countries.

May 2014 | VGU647308