

Drug Facts

Active ingredients (in each tablet)

Aspirin 500 mg (NSAID)*.....Pain reliever Caffeine 65 mg.....Alertness aid/pain reliever aid *nonsteroidal anti-inflammatory drug

Purpose

Uses

- for the temporary relief of minor aches and pains associated with a hangover
- helps restore mental alertness or wakefulness when experiencing fatigue or drowsiness associated with a hangover
- for temporary relief of headaches, body aches, and pain alone

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

hives • facial swelling • asthma (wheezing) • shock
 Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

• For occasional use only. Do not use for more than 2 days for a hangover unless directed by a doctor. Not intended for use as a substitute for sleep. If fatigue or drowsiness persists or continues to recur, consult a doctor.

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have a sodium-restricted diet



Alka-Seltzer Hangover Relief

Ask a doctor or pharmacist before use if you are taking a prescription drug for • diabetes • gout • arthritis		
 you experience a feel faint have stomach pain lasts for moder redness or swelli 	on occurs. Seek medical he ny of the following signs of omit blood • have bloody pain that does not get bette re than 10 days or gets wors ng is present s or a loss of hearing occurs	stomach bleeding: or black stools er se
use. It is especiall or later in pregnan doctor because it complications dur Keep out of reach help or contact a Po	est-feeding, ask a health pr y important not to use asy icy unless definitely direct may cause problems in th ing delivery. of children. In case of ove bison Control Center right av	pirin at 20 weeks ted to do so by a e unborn child or erdose, get medical
<i>Directions</i>fully dissolve 2 tag	blets in 4 ounces of water b	pefore taking
Adults 18 years and over	2 tablets every 6 hours as needed, or as directed by a doctor	do not exceed 8 tablets in 24 hours
Adults 60 years and over	2 tablets every 6 hours as needed, or as directed by a doctor	do not exceed 5 tablets in 24 hours
Children under 18 years	consult a doctor	
each tablet cont Phenylketonurid store at room ten Inactive ingredie dimethicone, docus	ains: potassium 115 mg ains: sodium 371 mg s: Contains Phenylalanine perature. Avoid excessive ants anhydrous citric acid, ate sodium, FD&C red No.	heat. aspartame, 40, FD&C yellow
No. 6, flavor, manni	tol, potassium bicarbonate, icarbonate, sucralose	
Questions or col 5PM EST)	mments? 1-800-986-0369	(Mon – Fri 9AM –